

Recommendations for psychological testing¹

Type of testing	Age range	Recommended test	Interval
Developmental and intellectual	<2.5 years	Bayley Scales of Infant and Toddler Development—Third Edition	As clinically indicated
	2.5-6 years	Wechsler Preschool and Primary Scale of Intelligence—Third Edition (WPPSI-III)	Every 3 years or as clinically indicated
	>6 years	Wechsler Abbreviated Scale of Intelligence—Second Edition (WASI-II)	As clinically indicated
Executive functioning	>3 years	Behavior Rating Inventory of Executive Functioning (BRIEF); self (age 11 and above), parent, and teacher report	At least every 2–3 years or as clinically indicated
Behavioral/emotional	3-18 years	Behavioral Assessment Scale for Children—Second Edition (BASC-2); self (age 8 and above), parent, and teacher report	At least every 2–3 years or as clinically indicated
	Adults	Beck Depression Inventory—Second Edition (BDI-II) and/or the Beck Anxiety Inventory (BAI)	At least every 2–3 years or as clinically indicated
Adaptive skills	>3 years	Adaptive Behavior Assessment System—Second Edition (ABAS-II)	At least every 2–3 years or as clinically indicated
	Adults	Adaptive Behavior Assessment System—Second Edition (ABAS-II)	As clinically indicated

¹Recommendations are derived from ACMG Practice Guidelines (F.2626; American College of Medical Genetics and Genomics Therapeutics Committee, Vockley, J., Andersson, H.C., Antshel, K.M., Braverman, N.E., Burton, B.K., Frazier, D.M., Mitchell, J., Smith, W.E., Thompson, B.H., & Berry, S.A. (Feb 2014). Phenylalanine hydroxylase deficiency: diagnosis and management guideline. *Genet. Med.*, 16, 188-200. doi:10.1038/gim.2013.157)